

Bone Graft

After Bone Graft

The following information applies when grafting material has been placed into extraction sites to help preserve your jawbone in preparation for possible implant replacement of the extracted tooth.

Your bone graft is made up of many particles. You may find some small granules in your mouth for the first several days. Do not be alarmed, it's normal to have some of them come out of the graft site and into your mouth. There are some things you could do to minimize the amount of particles that become dislodged:

- Do not disturb or touch the wound.
- Avoid rinsing or spitting for 2 days to allow blood clot and graft material stabilization.
- Do not apply pressure with your tongue or fingers to the grafted area, as the material is movable during the initial healing.
- Do not lift or pull on the lip to look at the sutures. This can actually cause damage to the wound site and tear the sutures.
- Do not smoke.

Rinsing

Following the second day, gentle rinsing is allowed but not too vigorously as you can again disturb some of the bone graft granules. If a partial denture or a flipper was placed in your mouth, you may have to see your restorative dentist to have it adjusted and learn how to remove and replace it appropriately.

Bleeding

When you leave our office after your bone graft there may be a special membrane placed on top of the bone graft and gauze placed above it. Try to keep the gauze in place until the evening. Once you remove the gauze you may feel some grains in your mouth. That is normal and to be expected, it does not mean the bone graft has been dislodged. Gently, with a tissue wipe the grains from your lips. If you have more than light bleeding after the first night please call our office.